

## GST Tag Antibody

Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP1298a

### Specification

#### GST Tag Antibody - Product Information

Application	WB,E
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig

#### GST Tag Antibody - Additional Information

##### Other Names

Green Fluorescent Protein

##### Target/Specificity

This GST Tag antibody is generated from rabbits immunized with a recombinant full length of GST.

##### Dilution

WB ~ 1:1000

##### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

##### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions


GST Tag Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### GST Tag Antibody - Protein Information

### GST Tag Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)



### GST Tag Antibody - Background

Glutathione S-transferase (GST) was originally cloned from parasite *Schistosoma japonicum* and it is now a widely used protein fusion partner. Vectors containing GST Tag have been developed for both prokaryotic and eukaryotic systems. The GST fusion proteins are easily purified from cell lysates by affinity chromatography using Glutathione Sepharose 4B to elute out the GST fusion protein from the column with a denaturing form of glutathione. Using the Abgent anti-GST antibody provides a simple solution to detect the expression of GST fusion proteins in cells.

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **GST Tag Antibody - Citations**

- [Mammalian sterile 20-like kinase 1/2 inhibit Wnt/ \$\beta\$ -catenin signaling pathway by directly binding Casein kinase 1 epsilon.](#)
- [Scd6 targets eIF4G to repress translation: RGG motif proteins as a class of eIF4G-binding proteins.](#)
- [Decapping activators in Saccharomyces cerevisiae act by multiple mechanisms.](#)