

LC3 Antibody (APG8A) (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1801A

Specification

LC3 Antibody (APG8A) (N-term) - Product Information

Application	WB, IF, IHC-P,E
Primary Accession	Q9H492
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Antigen Region	1-30

LC3 Antibody (APG8A) (N-term) - Additional Information

Gene ID 84557

Other Names

Microtubule-associated proteins 1A/1B light chain 3A, Autophagy-related protein LC3 A, Autophagy-related ubiquitin-like modifier LC3 A, MAP1 light chain 3-like protein 1, MAP1A/MAP1B light chain 3 A, MAP1A/MAP1B LC3 A, Microtubule-associated protein 1 light chain 3 alpha, MAP1LC3A

Target/Specificity

This LC3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1~30 amino acids from the N-term of human LC3 (APG8a).

Dilution

WB~~1000
IF~~1:200
IHC-P~~1:50~100

Format


Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage


Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LC3 Antibody (APG8A) (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.



Western blot analysis of anti-LC3 (APG8a) Pab (Cat. #AP1801a) in rat brain lysate. Both lipidated (arrow, II) and non-lipidated APG8a (arrow, I) were detected in membrane fraction (P) but only non-lipidated LC3 was detected in soluble fraction (S).



Fluorescent image of U251 cells stained with AP1801a LC3 (APG8A) (N-term) antibody. U251 cells were treated with Chloroquine (50 µM, 16h), then fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min). Cells were then incubated with AP1801a LC3 (APG8A) (N-term) primary antibody (1:200, 2 h at room temperature). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:1000, 1h). Nuclei were counterstained with Hoechst 33342 (blue) (10 µg/ml, 5 min). LC3 immunoreactivity is localized to autophagic vacuoles in the cytoplasm of U251 cells.

LC3 Antibody (APG8A) (N-term) - Protein Information

Name MAP1LC3A

Function

Ubiquitin-like modifier involved in formation of autophagosomal vacuoles (autophagosomes) (PubMed:20713600, PubMed:24290141). Whereas LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation (PubMed:20713600).

Cellular Location

Cytoplasm, cytoskeleton. Endomembrane system; Lipid-anchor. Cytoplasmic vesicle, autophagosome membrane; Lipid-anchor. Cytoplasmic vesicle, autophagosome
Note=LC3-II binds to the autophagic membranes


Tissue Location

Most abundant in heart, brain, liver, skeletal muscle and testis but absent in thymus and peripheral blood leukocytes.

LC3 Antibody (APG8A) (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)



Formalin-fixed and paraffin-embedded human brain tissue reacted with Autophagy LC3 Antibody (APG8a) (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

LC3 Antibody (APG8A) (N-term) - Background

Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. These proteins are involved in formation of autophagosomal vacuoles (autophagosomes). MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. MAP1LC3a is one of the light chain subunits and can associate with either MAP1A or MAP1B. The precursor molecule is cleaved by APG4B/ATG4B to form the cytosolic form, LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-bound form, LC3-II.

LC3 Antibody (APG8A) (N-term) - References

References for protein:

1. Baehrecke EH. Nat Rev Mol Cell Biol. 6(6):505-10. (2005)
2. Lum JJ, et al. Nat Rev Mol Cell Biol. 6(6):439-48. (2005)
3. Greenberg JT. Dev Cell. 8(6):799-801. (2005)
4. Levine B. Cell. 120(2):159-62. (2005)
5. Shintani T and Klionsky DJ. Science.

306(5698):990-5. (2004)

6.Tanida I., et al. Int. J. Biochem. Cell Biol. 36:2503-2518(2004)

7.He H., et al. J. Biol. Chem. 278:29278-29287(2003)

8.Tanida I., et al. J. Biol. Chem. 279:36268-36276(2004)

References for U251 cell line:

1. Westermark B.; Pontén J.; Hugosson R. (1973). "Determinants for the establishment of permanent tissue culture lines from human gliomas". Acta Pathol Microbiol Scand A. 81:791-805. [PMID: 4359449].

2. Pontén, J., Westermark B. (1978). "Properties of Human Malignant Glioma Cells in Vitro". Medical Biology 56: 184-193.[PMID: 359950].

3. Geng Y.; Kohli L.; Klocke B.J.; Roth K.A.(2010). "Chloroquine-induced autophagic vacuole accumulation and cell death in glioma cells is p53 independent". Neuro Oncol. 12(5): 473-481.[PMID: 20406898].

LC3 Antibody (APG8A) (N-term) - Citations

- [Baicalein Induces Autophagy and Apoptosis through AMPK Pathway in Human Glioma Cells.](#)
- [Induction of Autophagic Death of Human Hepatocellular Carcinoma Cells by Armillaridin from .](#)
- [Honokiol inhibits in vitro and in vivo growth of oral squamous cell carcinoma through induction of apoptosis, cell cycle arrest and autophagy.](#)
- [DOWNREGULATED APOPTOSIS AND AUTOPHAGY AFTER ANTI-A \$\beta\$ IMMUNOTHERAPY IN ALZHEIMER'S DISEASE.](#)
- [Procyanidins from Vitis vinifera seeds induce apoptotic and autophagic cell death via generation of reactive oxygen species in squamous cell carcinoma cells.](#)
- [Lmx1a and Lmx1b regulate mitochondrial functions and survival of adult midbrain dopaminergic neurons.](#)
- [Midazolam regulated caspase pathway, endoplasmic reticulum stress, autophagy, and cell cycle to induce apoptosis in MA-10 mouse Leydig tumor cells.](#)
- [Hyaluronan coated cerium oxide nanoparticles modulate CD44 and reactive oxygen species expression in human fibroblasts.](#)
- [A High Concentration of Genistein Induces Cell Death in Human Uterine Leiomyoma Cells by Autophagy.](#)
- [The role of tetraspanin CD9 in osteoarthritis using three different mouse models.](#)
- [Frontal white matter hyperintensities, clasmotodendrosis and gliovascular abnormalities in ageing and post-stroke dementia.](#)
- [Bach1 deficiency reduces severity of osteoarthritis through upregulation of heme oxygenase-1.](#)
- [Forkhead box protein o3 transcription factor negatively regulates autophagy in human cancer cells by inhibiting forkhead box protein o1 expression and cytosolic accumulation.](#)
- [Expression of LC3 and Beclin 1 in the spinal dorsal horn following spinal nerve ligation-induced neuropathic pain.](#)
- [Overexpression of the autophagic beclin-1 protein clears mutant ataxin-3 and alleviates Machado-Joseph disease.](#)
- [Identification and anti-human glioblastoma activity of tagitinin C from Tithonia diversifolia methanolic extract.](#)
- [IGF-I stimulates Rab7-RILP interaction during neuronal autophagy.](#)
- [The melanoma-associated transmembrane glycoprotein Gpnmb controls trafficking of cellular debris for degradation and is essential for tissue repair.](#)
- [Inhibition of isoprenylcysteine carboxylmethyltransferase induces autophagic-dependent apoptosis and impairs tumor growth.](#)
- [c-Abl and Arg tyrosine kinases regulate lysosomal degradation of the oncoprotein Galectin-3.](#)
- [Immunohistochemical evidence for macroautophagy in neurones and endothelial cells in Alzheimer's disease.](#)
- [Chloroquine-induced autophagic vacuole accumulation and cell death in glioma cells is p53 independent.](#)
- [Dual inhibition of akt/mammalian target of rapamycin pathway by nanoparticle albumin-bound-rapamycin and perifosine induces antitumor activity in multiple myeloma.](#)
- [Chondrocyte autophagy is stimulated by HIF-1 dependent AMPK activation and mTOR suppression.](#)
- [Heavy ion irradiation induces autophagy in irradiated C2C12 myoblasts and their bystander cells.](#)
- [Superoxide is the major reactive oxygen species regulating autophagy.](#)

- [Autophagy enhances the presentation of endogenous viral antigens on MHC class I molecules during HSV-1 infection.](#)
- [Autophagy protects neuron from Abeta-induced cytotoxicity.](#)