

Phospho-Bik(T33) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3042a

Specification

Phospho-Bik(T33) Antibody - Product Information

Application	WB, IHC-P,E
Primary Accession	Q13323
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	18016

Phospho-Bik(T33) Antibody - Additional Information

Gene ID 638

Other Names

Bcl-2-interacting killer, Apoptosis inducer NBK, BIP1, BP4, BIK, NBK

Target/Specificity

This Bik Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding T33 of human Bik.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-Bik(T33) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-Bik(T33) Antibody - Protein Information

Name BIK

Synonyms NBK

Function Accelerates programmed cell death. Association to the apoptosis repressors Bcl-X(L),

BHRF1, Bcl-2 or its adenovirus homolog E1B 19k protein suppresses this death-promoting activity. Does not interact with BAX.

Cellular Location


Endomembrane system; Single-pass membrane protein. Mitochondrion membrane; Single-pass membrane protein. Note=Around the nuclear envelope, and in cytoplasmic membranes

Phospho-Bik(T33) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-Bik(T33) Antibody - Images



Western blot analysis of anti-Bik Pab (Cat. #AP3042a) in mouse liver tissue lysate (35ug/lane). Bik(arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Phospho-Bik(T33) Antibody - Background

The protein encoded by this gene is known to interact with cellular and viral survival-promoting proteins, such as BCL2 and the Epstein-Barr virus in order to enhance programmed cell death. Because its activity is suppressed in the presence of survival-promoting proteins, this protein is suggested as a likely target for antiapoptotic proteins. This protein shares a critical BH3 domain with other death-promoting proteins, BAX and BAK.

Phospho-Bik(T33) Antibody - References

- Nikrad, M., et al., Mol. Cancer Ther. 4(3):443-449 (2005).
- Dong, F., et al., Infect. Immun. 73(3):1861-1864 (2005).
- Hur, J., et al., Proc. Natl. Acad. Sci. U.S.A. 101(8):2351-2356 (2004).
- Gillissen, B., et al., EMBO J. 22(14):3580-3590 (2003).
- Arena, V., et al., Genes Chromosomes Cancer 38(1):91-96 (2003).