

Anti-Bovine IgG (H&L) (Peroxidase Conjugated) Secondary Antibody
Rabbit Polyclonal, Peroxidase (Horseradish)
Catalog # ASR2018**Specification****Anti-Bovine IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Product Information**

Description	Anti-BOVINE IgG (RABBIT) Antibody
Host	Peroxidase Conjugated
Conjugate	Rabbit
Target Species	Peroxidase (Horseradish)
Clonality	Bovine
Application	Polyclonal
Application Note	,10,15, ELISA 1:10,000-1:50,000;Immunochemistry 1:500-1:2,500
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Bovine IgG whole molecule
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/ml Polyethylene Glycol (PEG-8000)
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Anti-Bovine IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

Anti-Bovine IgG antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Bovine IgG and Bovine Serum.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Bovine IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protein Information

Anti-Bovine IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Bovine IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Images

Anti-Bovine IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Background

Bovine IgG antibody detects heavy and light chain of Bovine IgG molecule. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Bovine IgG and Bovine Serum. Anti-Bovine IgG is ideal for investigators involved in serum protein components.