

# Anti-Cat IgG F(c) (Texas Red™ Conjugated) Secondary Antibody

Goat Polyclonal, Texas Red® Catalog # ASR2435

## **Specification**

## Anti-Cat IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Product Information

Description Anti-CAT IgG F(c) (GOAT) Antibody Texas

**Red™ Conjugated** 

Host Goat Conjugate Texas Red®

FP Value 3.1 moles Texas Red® per mole of IgG

Target Species Cat

Clonality Polyclonal Application ,3,4,

Application Note FLISA 1:10,000-1:50,000;IF Microscopy

1:1,000-1:5,000;FlowCytometry

1:500-1:2,500 Lyophilized

Physical State
Host Isotype
Target Isotype
IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Cat IgG F(c) fragment

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

**Immunoglobulin and Protease free** 

Preservative 0.01% (w/v) Sodium Azide

## Anti-Cat IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Additional Information

#### **Shipping Condition**

Ambient

Immunogen

#### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Cat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Cat IgG, Cat IgG F(c) and Cat Serum. No reaction was observed against Cat IgG F(ab')2

## **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.



## **Anti-Cat IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Protein Information**

# Anti-Cat IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Cat IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Images

Anti-Cat IgG F(c) (Texas Red™ Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.