

Anti-ARMENIAN HAMSTER IgG (H&L) (GOAT) (Peroxidase Conjugated) Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish)
 Catalog # ASR2512

Specification

Anti-ARMENIAN HAMSTER IgG (H&L) (GOAT) (Peroxidase Conjugated) Secondary Antibody - Product Information

Description	Anti-ARMENIAN HAMSTER IgG (H&L) (GOAT) Antibody Peroxidase Conjugated
Host	Goat
Conjugate	Peroxidase (Horseradish)
Target Species	Armenian Hamster
Reactivity	Hamster
Clonality	Polyclonal
Application	,1,10,15,
Application Note	ELISA 1:25,000;Western Blot 1:1,000-1:5,000;Immunochemistry 1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Armenian Hamster IgG whole molecule
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Anti-ARMENIAN HAMSTER IgG (H&L) (GOAT) (Peroxidase Conjugated) Secondary Antibody - Background

HRP Anti-Hamster IgG Secondary Antibody is designed for Western Blotting, ELISA and Immunohistochemistry. HRP conjugated secondary antibodies can also be used for a variety of other applications such as Assay Development.

Anti-ARMENIAN HAMSTER IgG (H&L) (GOAT) (Peroxidase Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Armenian Hamster IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Armenian Hamster IgG and Armenian Hamster Serum. Greatly diminished reactivity will occur against Golden Syrian Hamster IgG.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-ARMENIAN HAMSTER IgG (H&L) (GOAT) (Peroxidase Conjugated) Secondary Antibody - Protein Information**Anti-ARMENIAN HAMSTER IgG (H&L) (GOAT) (Peroxidase Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)