

MOUSE IgG3 isotype control Allophycocyanin
Monoclonal IgG3 , Allophycocyanine
Catalog # ASR3324**Specification**

MOUSE IgG3 isotype control Allophycocyanin - Product Information

Description	MOUSE IgG3 isotype control
Conjugate	Allophycocyanin Conjugated
FP Value	Allophycocyanine
Clonality	1-2 moles Allophycocyanine per mole of
Application	Mouse IgG3
Application Note	Monoclonal
Physical State	,4,10,
Host Isotype	ELISA 1:2000-1:20,000;FlowCytometry
Species of Origin	1:1000-1:5000
Stabilizer	Liquid (sterile filtered)
Preservative	IgG3
	Mouse
	None
	0.01% (w/v) Sodium Azide

MOUSE IgG3 isotype control Allophycocyanin - Additional Information**Shipping Condition**

Wet Ice

Purity

Mouse Isotype control has been prepared from concentrated cell culture supernatant by immunoaffinity chromatography using protein A. In an Ouchterlony double diffusion assay the material is non-reactive with antisera to mouse IgG1, IgG2a, IgG2b, IgM, and IgA. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Mouse IgG and anti-Mouse serum. Light and heavy chain composition has been confirmed.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. DO NOT FREEZE. This product is light sensitive.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

MOUSE IgG3 isotype control Allophycocyanin - Protein Information**MOUSE IgG3 isotype control Allophycocyanin - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MOUSE IgG3 isotype control Allophycocyanin - Images

MOUSE IgG3 isotype control Allophycocyanin - Background

Mouse isotype controls are used in flow cytometry, western blot and ELISA and differentiate between immunoglobulin classes and subclasses. Isotype controls allow for the genetic variations or differences in the constant regions of the heavy and light chains. In mouse there are six relevant heavy chain isotypes and two light chain isotypes: heavy chain α - IgA, γ - IgG 1, 2a, 2b, 3 and μ - IgM, light chain κ and λ .